



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. Solution by **M. C. STEVENS, M. A.**, Mathematical Department, Purdue University, Lafayette, Indiana; **HENRY HEATON, M. Sc.**, Atlantic, Iowa; **JOHN B. FAUGHT, A. M.**, Instructor in Mathematics in Indiana University, Bloomington, Indiana; and **J. C. GREGG, A. M.**, Brazil, Indiana.

Put $y=vx$ and we have

$$v + x \frac{dv}{dx} = \frac{vx^2 - v^2x^2}{x^2 + vx^2} = \frac{v - v^2}{1 + v} = v - \frac{2v^2}{1 + v}. \quad \text{Whence, } \frac{1+v}{v^2} dv + \frac{2dx}{x} = 0.$$

Integrating, $-1/v + \log(vx^2) + C = 0$, or $x/y = \log(xy) + C$, and $x = y \log(xy) + Cy$.

The C should not be omitted unless the conditions of the question giving rise to the equation are such as to make it zero.

IV. Solution by **H. C. WHITTAKER, A. M.**, Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

Let $y = x^p v^q$ and substitute in the given equation and we obtain

$$\frac{dv}{dx} = \frac{(1-p)v - (1+p)x^{p-1}v^{q+1}}{q^x(1+x^{p-1}v^q)}.$$

This will reduce to a simple form if we take $p=1$ and $q=-1$, giving

$$\frac{dv}{dx} = \frac{2}{x(1+v^{-1})}, \text{ or } dv(1+v^{-1}) = 2x^{-1}dx.$$

$$v + \log v = \log x^2; \quad x/y + \log(x/y) = \log x^2.$$

$$x/y = \log x^2 - \log(x/y) = \log(xy), \text{ whence } x = y \log(xy).$$

54. Proposed by **J. SCHEFFER, A. M.**, Hagerstown, Maryland.

A certain solid has a square, side= a , for its base, and all parallel sections are squares, the two sections through the middle points of the opposite side of the square are semi-circles, however. Find surface, volume, and center of gravity of each.

I. Solution by **HENRY HEATON, M. Sc.**, Atlantic, Iowa.

The length of a side of a parallel section distant x from the base is $(a^2 - 4x^2)^{\frac{1}{2}}$. If dx be the distance between two parallel sections, the distance between two corresponding sides is $adx/(a^2 - 4x^2)^{\frac{1}{2}}$. Hence the surface

$$S = 4 \int_0^{\frac{1}{2}a} adx = 2a^2; \text{ the volume } V = \int_0^{\frac{1}{2}a} (a^2 - 4x^2)dx = \frac{1}{3}a^3;$$

the distance of the center of gravity of the surface from the base is

$$\frac{1}{2a^2} \int_0^{\frac{1}{2}a} axdx = \frac{1}{6}a;$$

and the distance of the center of gravity of the volume from the base is

$$\frac{3}{a^3} \int_0^{\frac{1}{2}a} x(a^2 - 4x^2) dx = \frac{3}{16}a.$$

II. Solution by J. C. NAGLE, M. A., M. C. E., Professor of Civil Engineering in the State A. M. College, College Station, Texas.

Take the intersection of the planes of the circular sections as the axis of z , the origin being in the center of the base. Then since the radius of each circle is $\frac{1}{2}a$ we shall have for the projection of one fourth of the elementary area intercepted between two planes parallel to the base, and distant dz from each other, upon the plane of one of the circles,

$$dS \cos \theta = \sqrt{\frac{1}{4}a^2 - z^2} \cdot dz,$$

where θ is the angle made by this elementary area with the plane of projection.

$$\text{But } \cos \theta = \frac{\sqrt{\frac{1}{4}a^2 - z^2}}{\frac{1}{2}a}, \text{ and the whole surface is } S = 4 \int_0^{\frac{1}{2}a} a dz = 2a^2 \dots (1).$$

The center of gravity of S is distant from the base

$$z_1 = \frac{4 \int_0^{\frac{1}{2}a} a z dz}{2a^2} = \frac{1}{4}a \dots \dots \dots (2).$$

For the volume, taking planes parallel to the base,

$$V = \int_0^{\frac{1}{2}a} 2 \sqrt{\frac{1}{4}a^2 - z^2} \cdot 2 \sqrt{\frac{1}{4}a^2 - z^2} \cdot dz = \int_0^{\frac{1}{2}a} (a^2 - 4z^2) dz = a^3/3 \dots \dots \dots (3),$$

and its center of gravity above the base is

$$z_2 = \frac{\int_0^{\frac{1}{2}a} (a^2 - 4z^2) z dz}{\frac{1}{3}a^3} = \frac{3}{16}a \dots \dots \dots (4).$$

The figure will be a cloistered arch formed by the intersection of two right semi-cylinders.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

Let $x^2 + z^2 = \frac{1}{4}a^2 \dots \dots (1)$, $y^2 + z^2 = \frac{1}{4}a^2 \dots \dots (2)$ be the equations to the cylinders which form the groin. From (1) $dz/dx = -x/z$, $dz/dy = 0$.

$$\begin{aligned}
 S &= \iint \sqrt{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2} dx dy = 8 \int_0^a \int_0^x \sqrt{1 + \frac{x^2}{z^2}} dx dy \\
 &= 4a \int_0^a \int_0^x \frac{dx dy}{z} = 4a \int_0^a \int_0^x \frac{dx dy}{\sqrt{\frac{1}{4}a^2 - x^2}} = 4a \int_0^a \frac{xdx}{\sqrt{\frac{1}{4}a^2 - x^2}} = 2a^2.
 \end{aligned}$$

$$V = \iiint dz dx dy = 4 \int_0^a \int_0^{\sqrt{\frac{1}{4}a^2 - 4z^2}} \int_0^{\sqrt{\frac{1}{4}a^2 - 4z^2}} dz dx dy = \int_0^a (a^2 - 4z^2) dz = \frac{1}{3}a^3.$$


$$\text{Center of gravity of surface} = \frac{\iint z dS}{\iint dS} = \frac{1}{2}a \int_0^a \int_0^x dx dy = \frac{1}{2}a.$$

$$\text{Center of gravity of volume} = \frac{\iiint z dz dx dy}{\iiint dz dx dy} = \frac{\frac{3}{a^3} \int_0^a z(a^2 - 4z^2) dz}{\frac{1}{3}a^3} = \frac{3}{8}a.$$

IV. Solution by J. C. GREGG, A. M., Brazil, Indiana.

Let the given figure represent a section of the solid through the middle point of two opposite sides of the base. We have $r = a/2$, and the equation of the circle EDF is $x^2 + y^2 = r^2 \dots (1)$, and $AC^2 = (2y)^2 = 4(r^2 - x^2) = A_x = a$ section parallel to the base, and for the volume

$$V = 4 \int_0^r (r^2 - x^2) dx = \frac{8}{3}r^3 = \frac{1}{3}a^3.$$



The surface may be considered to be generated by the sides of a section parallel to the base, and we have for the surface,

$$S = 4 \int 2y ds = 4 \int_0^r 2\sqrt{r^2 - x^2} \cdot \frac{r dx}{\sqrt{r^2 - x^2}} = 8r \int_0^r dx = 8r^2 = 2a^2.$$

For the center of gravity of the volume,

$$\bar{x} = \frac{\int x(A_x) dx}{V} = \frac{4 \int_0^r x(r^2 - x^2) dx}{\frac{8}{3}r^3} = \frac{3}{2r^3} \int_0^r x(r^2 - x^2) dx = \frac{3}{8}r = \frac{3}{8}a.$$

For the center of gravity of the *curved* surface we have,

$$\bar{x} = \frac{4 \int xy ds}{S}, = \frac{8r \int_0^r x dx}{8r^2} = \frac{1}{r} \int_0^r x dx, = \frac{1}{2}r, = \frac{1}{2}a.$$

For the center of gravity of the *whole* surface, since the curved surface is *twice* that of the base we have, $\bar{x} = \frac{2}{3} \cdot \frac{1}{2}a = \frac{1}{3}a$.

Also solved by H. C. WHITAKER, C. W. M. BLACK, and the PROPOSER.

Professors Black and Scheffer used "side= $2a$ " as in Problem 47, instead of *side*= a , and hence their results did not agree with those in the published solutions. The results obtained were : Volume= $8a^3/3$, surface= $8a^2$, center of gravity of volume= $3a/8$, and center of gravity of surface= $\frac{1}{2}a$. See problem 42 for two additional SOLUTIONS for surface and volume. EDITOR.

MECHANICS.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

38. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

A prolate spheroid of revolution is fixed at its focus ; a blow is given it at the extremity of the axis minor in a line tangent to the direction perpendicular to the axis major. Find the axis about which the body begins to rotate. [From *Loudon's Rigid Dynamics*.]

Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

The general equations of motion are :

$$\left. \begin{aligned} A\omega_x - (\Sigma mxy)\omega_y - (\Sigma mxz)\omega_z &= L \\ B\omega_y - (\Sigma myz)\omega_z - (\Sigma myx)\omega_x &= M \\ C\omega_z - (\Sigma mzx)\omega_x - (\Sigma mzy)\omega_y &= N \end{aligned} \right\} \dots\dots\dots (1).$$

The equation to the ellipsoid with focus as origin is $a^2y^2 + a^2z^2 + b^2x^2 = 2aeb^2x + b^4$. Now $\Sigma mxy = \Sigma mxz = \Sigma myz = 0$. \therefore (1) reduce to

$$\left. \begin{aligned} A\omega_x &= L \\ B\omega_y &= M \\ C\omega_z &= N \end{aligned} \right\} \dots\dots\dots (2).$$